

MYCOTAXON

<http://dx.doi.org/10.5248/120.81>

Volume 120, pp. 81–88

April–June 2012

Two new freshwater species of *Annulatascaceae* from ChinaDIAN-MING HU^{1,2}, LEI CAI^{1*}, ALI HASSAN BAHKALI³ & KEVIN D. HYDE^{2,3,4,5*}¹State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing 100101, China²International Fungal Research & Development Centre, The Research Institute of Resource Insects, Chinese Academy of Forestry, Bailongsi, Kunming 650224, China³Botany and Microbiology Department, College of Science, King Saud University, Riyadh, Saudi Arabia⁴School of Science, Mae Fah Luang University, Chiang Rai, Thailand⁵Institute of Excellence for Fungal Research, Mae Fah Luang University, Chiang Rai, Thailand* CORRESPONDENCE TO: mrcailei@gmail.com, kdhyde3@gmail.com

ABSTRACT — Two new species belonging to *Annulatascaceae* are described and illustrated on submerged wood from a freshwater stream. *Annulatascus menglensis*, differentiated from other *Annulatascus* species by its hyaline ascomatal neck, is also characterized by cylindrical asci with a large bipartite refractive J– apical ring and hyaline fusiform ascospores without sheath or appendages. *Aqualignicola vaginata*, which is distinguished from *A. hyalina* in producing a gelatinous sheath surrounding the unicellular hyaline ascospores, is further characterized by its membranous ascomata with lanceolate setae that surround the neck, long cylindrical asci with a relatively large apical ring.

KEY WORDS — aquatic, *Ascomycota*, fungi, taxonomy

Introduction

Freshwater fungi are an important ecological group because of their indispensable function as decomposers in freshwater ecosystems (Wong et al. 1998a); they also have potential application in producing various and useful bioactive secondary metabolites (Bucher et al. 2004, Zhang et al. 2008, Hosoe et al. 2010, Dong et al. 2011). Freshwater fungi are however, relatively poorly investigated comparing to their terrestrial counterpart. Until recently, freshwater fungi had been reported mainly from Europe, North America, and Southeast Asia (Cai et al. 2006b, Raja et al. 2009). Although China has a high biodiversity of fungi and abundant freshwater resources, investigation of freshwater fungi has been mostly restricted to Hong Kong and Yunnan Province.

Annulatascaceae S.W. Wong et al. comprises many freshwater genera (Vijaykrishna et al. 2006). The family is characterized morphologically by dark

brown or black ascomata with ostiolate necks, long cylindrical unitunicate asci with a relatively massive, refractive apical ring, and uniseriate, hyaline to brown, fusiform or ellipsoidal ascospores (Wong et al. 1998b, Ho & Hyde 2000). Seventy-five species representing 21 genera are currently placed in *Annulatascaceae* (Kirk et al. 2008).

Our research group has focused on investigating the diversity of freshwater fungi in China for over ten years (Hyde et al. 2000, Luo et al. 2004, Cai et al. 2005, 2008, Hu et al. 2007, 2010a, b). During our continuing investigation, two undescribed ascomycetes belonging to the *Annulatascaceae* were collected on submerged wood and are described and illustrated in this paper.

Materials & methods

D.M. Hu, who has investigated freshwater ascomycetes on submerged woody material in the southern provinces of China (including Guizhou, Jiangxi and Yunnan provinces) since 2008, has collected 875 samples from streams, lakes, ditches, and ponds. After return to the laboratory, the samples were incubated following Cai et al. (2006a). Samples were examined for fungal fruiting bodies under a dissecting microscope (Leica MZ16A) during incubation. Observations and photographs were prepared from materials mounted in water under a compound microscope (Nikon E800). Single spore isolation as outlined by Choi et al. (1999) was attempted but unsuccessful. Herbarium specimens of the fungal taxa described in this study are deposited in the International Fungi Research and Development Centre (IFRDC).

Taxonomy

Annulatascus menglensis D.M. Hu, L. Cai & K.D. Hyde, sp. nov.

PLATE 1

MYCOBANK MB 563810

Differs from other *Annulatascus* species by its hyaline neck.

TYPE: China, Yunnan Province, Mengla, Wudaoban Stream, 21°24'N 101°36'E, alt: 660 m, 3 April 2009, D.M. Hu (*Holotype*, IFRDC 023-002).

ETYMOLOGY: *menglensis* refers to the collecting site, Mengla, a county in southern China.

ASCOMATA 180–200 high, 120–150 µm in diam., subglobose to ellipsoidal, superficial, solitary, ostiolate, black, coriaceous. NECK 150–340 × 40–60 µm, cylindrical, hyaline, membranous, periphysate. PERIDIUM composed of compressed cells, outer layers dark brown, inner layers hyaline. PARAPHYSES ca. 5–6 µm wide, broad at the base and tapering towards the apex, hyaline, septate, smooth-walled, simple, longer than asci. ASCI 115–163 × 10–13 µm, 8-spored, cylindrical, unitunicate, persistent, pedicellate, with a large bipartite, refractive, J- apical ring, ca. 3 µm high, 4 µm in diam. ASCOSPORES 21–25 × 8–10 µm (mean = 23.8 × 8.5 µm, n = 30), uniseriate, overlapping, fusiform, with acute ends, hyaline, aseptate, guttulate, smooth, thin-walled, without sheaths or appendages.



PLATE 1. *Annulatascus menglensis* (from holotype). a. Ascoma on submerged wood. b. Squash mount of an ascoma. c. Peridium comprising angular cells in surface view. d. Paraphyses. e–f. Asci. g. Upper part of an ascus showing the apical ring. h–k. Ascospores. Scale bars: a = 100 μ m, b = 50 μ m, c–d = 5 μ m, e–f = 20 μ m, g–k = 5 μ m.

HABITAT & DISTRIBUTION— Saprobic on submerged wood in fresh water in China.

ADDITIONAL SPECIMENS EXAMINED: CHINA. YUNNAN PROVINCE: Mengla, Wudaoban Stream, 21°24'N 101°36'E, alt. 660 m, 3 April 2009, D.M. Hu (IFRDSC 002-017, IFRDSC 002-018).

COMMENTS — *Annulatascus menglensis* has black, coriaceous ascomata and cylindrical asci with a large bipartite, refractive, J– apical ring. It fits well within the generic concept of *Annulatascus* (Hyde 1992). Thus far, 14 species have been included in the genus (Tsui et al. 2002, Barbosa et al. 2008, Abdel-Wahab et al. 2011). The new species resembles *A. velatisporus* K.D. Hyde (Hyde 1992) in having fusiform and aseptate ascospores. However, *A. velatisporus* differs from *A. menglensis* in having larger ascospores ($26\text{--}42 \times 9\text{--}12\ \mu\text{m}$) with a gelatinous sheath. *Annulatascus menglensis* shares similar ascospore dimensions with *A. fusiformis* K.D. Hyde & S.W. Wong (Hyde & Wong 2000), *A. joannae* K.M. Tsui et al. (Tsui et al. 2002), *A. citriosporus* J. Fröhl. & K.D. Hyde (Fröhlich & Hyde 2000), and *A. apiculatus* F.R. Barbosa & Gusmão (Barbosa et al. 2008) but lacks their black necks and mucilaginous sheaths and/or appendages. *Annulatascus lacteus* K.M. Tsui et al. (Tsui et al. 2002), which also has aseptate ascospores without sheath or appendage, differs from *A. menglensis* in having a milky ascomata.

Aqualignicola vaginata D.M. Hu, L. Cai & K.D. Hyde, **sp. nov.**

PLATE 2

MYCOBANK MB 563811

Differs from *Aqualignicola hyalina* in having a hyaline ascomatal neck and a gelatinous sheath surrounding the ascospores.

TYPE: China, Yunnan Province, Mengla, Wudaoban Stream, 21°32'N E101°29'E, alt. 620 m, 2 April 2009, D.M. Hu (**Holotype**, IFRDC 021-043).

ETYMOLOGY: *vaginata* (= sheath) refers to the mucilaginous sheath surrounding the ascospore.

ASCOMATA 100–140 μm diam, globose, membranous, brown to black-brown, submerged, solitary. **NECK** 40–60 μm diam, 200–350 μm long, cylindrical, hyaline, periphysate, covered with setae; setae $15\text{--}35 \times 4\text{--}5\ \mu\text{m}$, lanceolate, solid, surrounding the neck. **PERIDIUM** 7–14 μm thick, in surface view of *textura angularis*, composed of 5 layers of light-brown, pseudoparenchymatous cells. **PARAPHYSES** 4–6 μm in diam, 85–100 long, septate, flexuose, hypha-like, numerous, tapering towards the apex, embedded in a gelatinous matrix. **ASCI** $145\text{--}156 \times 5.5\text{--}6.5\ \mu\text{m}$, 8-spored, cylindrical, pedicellate, unitunicate, apically truncate, with a large, J–, refractive, apical ring. **ASCOSPORES** $11\text{--}15 \times 5\text{--}6\ \mu\text{m}$ (mean = $13.4 \times 5.3\ \mu\text{m}$, $n = 50$), uniseriate, ellipsoidal-fusiform, hyaline, aseptate, thin-walled, multiguttulate, with a fan-shaped, hyaline mucilaginous sheath surrounding one end of the ascospores.




PLATE 2. *Aqualignicola vaginata* (from holotype). a. Ascoma on submerged wood. b. Section of an ascoma. c. Peridium. d. Section through a neck. e. Hair on neck. f. Paraphyses. g. Ascus. h. Upper part of an ascus, show the apical ring. i–n. Ascospores (arrows show the mucilaginous sheath and the multiple guttules). Scale bars: a = 100 μm , b = 30 μm , c = 5 μm , d = 30 μm , e, f = 5 μm , g = 20 μm , h–n = 5 μm .

HABITAT & DISTRIBUTION — Saprobic on submerged wood in fresh water in China.

ADDITIONAL SPECIMENS EXAMINED: CHINA. YUNNAN PROVINCE: Mengla, Wudaoban Stream, 21°32'N 101°29'E, alt. 620 m, 2 April 2009, D.M. Hu (IFRDSC 004-003, IFRDSC 004-004, IFRDSC 004-005).

COMMENTS — Previously monotypic, *Aqualignicola* V.M. Ranghoo et al. is characterized by membranous ascomata with a neck surrounded by lanceolate setae, long cylindrical asci with a relatively large apical ring, and hyaline unicellular ascospores (Ranghoo et al. 2001). The new species fits well with the generic concept of *Aqualignicola* but differs from the type species, *A. hyalina*, in having a hyaline ascomatal neck and a mucilaginous sheath surrounding the ascospore. Furthermore, the ascospores and asci of *A. vaginata* are narrower than those of *A. hyalina* (ascospores: 14–15 × 6.25–7.5 µm; asci 9–12 µm diam.). *Aqualignicola vaginata* is also similar to species of *Aquaticola* W.H. Ho et al. (Ho et al. 1999) and *Longicollum* Zelski et al. (Zelski et al. 2011a). However, the species of the latter two genera have glabrous ascomata and lack the lanceolate setae surrounding the neck found in *A. vaginata*. *Aqualignicola vaginata* can be compared to the newly described *Chaetorostrum quincemilense* Zelski et al. (Zelski et al. 2011b) in that both taxa have a hyaline neck covered with setae; however the ascospores of *A. vaginata* are aseptate and much smaller than those of *C. quincemilense* (1–3-septate; 30–38 × 10–12 µm).

Acknowledgments

Funds for research were provided by a major project of the National Natural Science Foundation of China, No. 31093440. The Global Research Network for Fungal Biology and King Saud University are thanked for support.

Literature cited

- Abdel-Wahab MA, Abdel-Aziz FA, Mohamed SS, Abdel-Aziz AE. 2011. *Annulatascus nilensis* sp. nov., a new freshwater ascomycete from the River Nile, Egypt. IMA Fungus 2: 1–6. <http://dx.doi.org/10.5598/ima fungus.2011.02.01.01>
- Barbosa FR, Gusmao LFP, Raja HA, Shearer CA. 2008. *Annulatascus apiculatus* sp. nov., a new freshwater ascomycete from the semi-arid Caatinga Biome of Brazil. Mycotaxon 106: 403–407.
- Bucher VVC, Pointing SB, Hyde KD, Reddy CA. 2004. Production of wood decay enzymes, loss of mass, and lignin solubilization in wood by diverse tropical freshwater fungi. Microbial Ecology 48: 331–337. <http://dx.doi.org/10.1007/s00248-003-0132-x>
- Cai L, Zhang KQ, Hyde KD. 2005. *Ascoyunnania aquatica* gen. et sp. nov., a freshwater fungus collected from China and its microcyclic conidiation. Fungal Diversity 18: 1–8.
- Cai L, Ji KF, Hyde KD. 2006a. Variation between freshwater and terrestrial fungal communities on decaying bamboo culms. Antonie Van Leeuwenhoek 89: 293–301. <http://dx.doi.org/10.1007/s10482-005-9030-1>
- Cai L, Hyde KD, Tsui CKM. 2006b. Genera of freshwater fungi. Fungal Diversity Research Series 18: 1–261.

- Cai L, Guo XY, Hyde KD. 2008. Morphological and molecular characterisation of a new anamorphic genus *Cheirosporium*, from freshwater in China. *Persoonia* 20: 53–58.
<http://dx.doi.org/10.3767/003158508X314732>
- Choi YW, Hyde KD, Ho WH. 1999. Single spore isolation of fungi. *Fungal Diversity* 3: 29–38.
- Dong JY, Shen KZ, Sun R. 2011. Recent progress in the research on bioactive substances from freshwater fungi. *Mycosystema* 30: 206–217.
- Fröhlich J, Hyde KD (eds). 2000. *Palm microfungi*. Fungal Diversity Press, Hong Kong.
- Ho WH, Hyde KD. 2000. A new family of freshwater ascomycetes. *Fungal Diversity* 4: 21–36.
- Ho W, Tsui CKM, Hodgkiss IJ, Hyde KD. 1999. *Aquaticola*, a new genus of *Annulatascaceae* from freshwater habitats. *Fungal Diversity* 3: 87–97.
- Hosoe T, Gloer JB, Wicklow DT, Raja HA, Shearer CA. 2010. New nonadride analogues from a freshwater isolate of an undescribed fungus belonging to the order *Pleosporales*. *Heterocycles* 81: <http://dx.doi.org/10.3987/COM-10-12009>
- Hu DM, Zhu H, Cai L, Hyde KD, Zhang KQ. 2007. *Sirothecium triseriale*, a new chirosporous anamorphic species from China. *Cryptogamie Mycologie* 28: 311–314.
- Hu DM, Cai L, Chen H, Bahkali AH, Hyde KD. 2010a. Four new freshwater fungi associated with submerged wood from Southwest Asia. *Sydowia* 62: 191–203.
- Hu DM, Cai L, Chen H, Bahkali AH, Hyde KD. 2010b. Fungal diversity on submerged wood in a tropical stream and an artificial lake. *Biodiversity and Conservation* 19: 3799–3808.
<http://dx.doi.org/10.1007/s10531-010-9927-5>
- Hyde KD. 1992. Tropical Australian freshwater fungi II. *Annulatascus velatispora* gen. et sp. nov. *Annulatascus bipolaris* sp. nov. and *Nais aquatica* sp. nov. *Australian Systematic Botany* 5: 117–124. <http://dx.doi.org/10.1071/SB9920117>
- Hyde KD, Wong SW. 2000. *Annulatascus fusiformis* sp. nov., a new freshwater ascomycete from the Philippines. *Mycologia* 92: 553–557. <http://dx.doi.org/10.2307/3761515>
- Hyde KD, Ho WH, Jones EBG, Tsui CKM, Wong WSW. 2000. *Torrentispora fibrosa* gen. sp. nov. (*Annulatascaceae*) from freshwater habitats. *Mycological Research* 104: 1399–1403.
<http://dx.doi.org/10.1017/S0953756200002781>
- Kirk PM, Cannon PF, Minter DW, Stalpers JA. 2008. *Dictionary of the fungi*, 10th edition. Cromwell Press, Trowbridge. 771 p.
- Luo J, Yin JF, Cai L, Zhang KQ, Hyde KD. 2004. Freshwater fungi in Lake Dianchi, a heavily polluted lake in Yunnan, China. *Fungal Diversity* 16: 93–112.
- Raja HA, Schmit JP, Shearer CA. 2009. Latitudinal, habitat and substrate distribution patterns of freshwater ascomycetes in the Florida Peninsula. *Biodiversity and Conservation* 18: 419–455.
<http://dx.doi.org/10.1007/s10531-008-9500-7>
- Ranghoo VM, Tsui CKM, Hyde KD. 2001. *Brunneosporella aquatica* gen. et sp. nov., *Aqualignicola hyalina* gen. et sp. nov., *Jobellisia viridifusca* sp. nov. and *Porosphaerellopsis bipolaris* sp. nov. (ascomycetes) from submerged wood in freshwater habitats. *Mycological Research* 105: 625–633. <http://dx.doi.org/10.1017/S0953756201003793>
- Tsui CKM, Ranghoo VM, Hodgkiss IJ, Hyde KD. 2002. Three new species of *Annulatascus* (*Ascomycetes*) from Hong Kong freshwater habitats. *Mycoscience* 43: 383–389.
<http://dx.doi.org/10.1007/s102670200056>
- Vijaykrishna D, Jeewon R, Hyde KD. 2006. Molecular taxonomy, origins and evolution of freshwater ascomycetes. *Fungal Diversity* 23: 351–390.
- Wong MKM, Goh TK, Hodgkiss IJ, Hyde KD, Ranghoo VM, Tsui CKM, Ho WH, Wong WSW, Yuen TK. 1998a. Role of fungi in freshwater ecosystems. *Biodiversity and Conservation* 7: 1187–1206. <http://dx.doi.org/10.1023/A:1008883716975>

- Wong SW, Hyde KD, Jones EBB. 1998b. *Annulatascaceae*, a new ascomycete family from the tropics. *Systema Ascomycetum* 16: 17–25.
- Zhang KQ, Wang L, Dong JY, Song HC, Shen KZ, Wang LM, Sun R, Wang CR, Li GH, Li L. 2008. Screening and isolation of antibacterial activities of the fermentative extracts of freshwater fungi from Yunnan Province, China. *Annals of Microbiology* 58: 579–584.
<http://dx.doi.org/10.1007/BF03175561>
- Zelski SE, Raja HA, Miller AN, Barbosa FR, Gusmao LFP, Shearer CA. 2011a. *Longicollum biappendiculatum* gen. et sp. nov., a new freshwater ascomycete from the Neotropics. *Mycosphere* 2: 539–545.
- Zelski SE, Raja HA, Miller AN, Shearer CA. 2011b. *Chaetorostrum quincemilensis*, gen. et sp. nov., a new freshwater ascomycete and its *Taeniolella*-like anamorph from Peru. *Mycosphere* 2: 593–600.